

Corporate Overview Presentation

February 2026

Legal disclaimer

Forward-Looking Statements

Statements in this presentation that are not statements of historical fact are forward-looking statements. Such forward-looking statements include, without limitation, statements regarding: the therapeutic and commercial potential of Vir Biotechnology's CHD program, as well as Vir Biotechnology's strategy, plans and expectations related thereto; the therapeutic and commercial potential of Vir Biotechnology's oncology solid tumor portfolio, preclinical pipeline and the PRO-XTEN® masking technology, as well as Vir Biotechnology's strategy, plans and expectations related thereto; the potential of and Vir Biotechnology's expectations for its other pipeline programs; the potential benefits for Vir Biotechnology as a result of the collaborations with Norgine and Astellas; Vir Biotechnology's anticipated cash runway; Vir Biotechnology's plans and expectations for its clinical development programs, including protocols for and enrollment into ongoing and planned clinical studies, potential partnering opportunities, and data readouts and presentations, as well as anticipated timelines; the potential benefits, safety and efficacy of Vir Biotechnology's investigational therapies; and any assumptions underlying any of the foregoing. Words such as “aim,” “anticipate,” “believe,” “could,” “expect,” “goal,” “intend,” “may,” “plan,” “potential,” “promising,” “will,” and similar expressions are intended to identify forward-looking statements, though not all forward-looking statements necessarily contain these identifying words. These forward-looking statements are based on the beliefs of the management of Vir Biotechnology, as well as assumptions made by and information currently available to management. Such statements reflect the current views of Vir Biotechnology with respect to future events and are subject to known and unknown risks, including, without limitation: unexpected safety or efficacy data or results observed during clinical studies or in data readouts, including the occurrence of adverse safety events; risks of unexpected costs, delays or other unexpected hurdles; the timing and amount of Vir Biotechnology's actual operating expenses, as determined in accordance with U.S. Generally Accepted Accounting Principles; difficulties in collaborating with other companies, some of whom may be competitors of Vir Biotechnology or otherwise have divergent interests, and uncertainty as to whether the benefits of Vir Biotechnology's various collaborations can ultimately be achieved; challenges in accessing manufacturing capacity; clinical site activation rates or clinical enrollment rates that are lower than expected; the timing and outcome of Vir Biotechnology's planned interactions with regulatory authorities, as well as general difficulties in obtaining any necessary regulatory approvals; successful development and/or commercialization of alternative product candidates by Vir Biotechnology's competitors, as well as changes in expected or existing competition; Vir Biotechnology's use of AI and machine learning in its efforts to engineer next-generation proteins and in other research and development efforts; geopolitical changes or other external factors; and unexpected litigation or other disputes. In light of these risks and uncertainties, the events or circumstances referred to in the forward-looking statements may not occur. Drug development and commercialization involve a high degree of risk, and only a small number of research and development programs result in commercialization of a product. Results in early-stage clinical studies may not be indicative of full results or results from later stage or larger scale clinical studies and do not ensure regulatory approval. The actual results may vary from the anticipated results and the variations may be material. Other factors that may cause Vir Biotechnology's actual results to differ from current expectations are discussed in Vir Biotechnology's filings with the U.S. Securities and Exchange Commission, including the section titled “Risk Factors” contained therein. These forward-looking statements should not be taken as forecasts or promises nor should they be taken as implying any indication, assurance or guarantee that the assumptions on which such forward-looking statements have been made are correct or exhaustive or, in the case of the assumptions, fully stated in this presentation. You are cautioned not to place undue reliance on any scientific data presented or these forward-looking statements, which speak only as of the date of this presentation. Except as required by law, Vir Biotechnology undertakes no obligation to publicly update any forward-looking statements, whether as a result of new information, future events or otherwise. Vir Biotechnology claims the protection of the safe harbor for forward-looking statements contained in the Private Securities Litigation Reform Act of 1995 for all forward-looking statements.

Product candidates included in this presentation are investigational and have not been approved by the US Food and Drug Administration or other regulatory authorities. No representation is made or intended regarding their safety or efficacy or that of other investigational agents mentioned herein. Any comparative data presented are based on cross-trial comparisons and not head-to-head clinical studies; therefore, caution should be exercised in interpreting these data.



POWERING THE
IMMUNE SYSTEM TO
TRANSFORM
LIVES

Our path to delivering transformational therapies to people living with devastating diseases



Commercializing our **chronic hepatitis delta (CHD)** combination therapy will drive near-term revenue sustainability



Accelerating our **masked T-cell engager (TCE) immunotherapy** portfolio offers key value inflection points



Aiming Vir Bio's discovery engine at **developing a robust pipeline** of cancer immunotherapies creates sustainable long-term growth



Strategic Collaborations

Selectively partner drug candidates to focus internal resources, unlock the value of our pipeline and maximize benefit to patients

We've developed a powerful Vir Bio discovery engine to fuel the next generation of therapeutics

Our distinctive capabilities



**World class protein engineering,
antibody / TCE discovery**



**dAlsY™ AI/ML for antibody / TCE
optimization**



**Exclusive PRO-XTEN® universal
masking technology**

**Building on legacy of
infectious disease innovation**

Ebanga™

(ansuvimab-zykl)

for the treatment of
ebola virus

Xevudy®

(sotrovimab)

for the treatment of
SARS-COVID 19

**to deliver next generation of powerful
medicines, including cancer
immunotherapies with better therapeutic
index**

Delivering a differentiated pipeline in oncology and infectious disease

Driving near-term and long-term value creation



Disease Area	Product Candidate	Goal	Pre-clinical	Phase 1	Phase 2	Phase 3	Approval
CLINICAL PROGRAMS							
Chronic Hepatitis Delta	tobevibart + elebsiran	Treatment					
Solid Tumors	VIR-5500 (PSMA) ¹ ± ARPIs	Treatment					
Solid Tumors	VIR-5818 (HER2) ¹ ± pembrolizumab	Treatment					
Solid Tumors	VIR-5525 (EGFR) ¹ ± pembrolizumab	Treatment					
PRE-CLINICAL PROGRAMS							
HIV Treatment / Cure ²	Preclinical antibody candidates	Treatment					
Solid Tumors	7 PRO-XTEN® TCE programs including lung, colorectal and bladder cancers	Treatment					

¹ Masked TCEs licensed from Sanofi

² In collaboration with the Gates Foundation

ARPIs: androgen receptor pathway inhibitors; EGFR: epidermal growth factor receptor; HER2: human epidermal growth factor receptor 2; HIV: human immunodeficiency virus; PSMA: prostate-specific membrane antigen; siRNA: small interfering RNA; TCE: T-cell engager

Tobevibart incorporates Xencor's Xtend[™] and other Fc technologies

Norgine holds exclusive license for the commercial rights to the combination of tobevibart and elebsiran in Europe, Australia and New Zealand

Brii Biosciences retains rights to the combination of tobevibart and elebsiran in the Greater China Territory (People's Republic of China, Hong Kong, Taiwan and Macau)

Upon closing of the Astellas global collaboration, which is contingent on customary closing conditions, including clearance under the Hart-Scott-Rodino (HSR) Act, Astellas will hold co-development and co-commercialization rights for VIR-5500 for the treatment of prostate cancer.

Upcoming clinical milestones

PROGRAM	DRUG CANDIDATES REGIMEN	CATALYST	TIMING
Hepatitis Delta	tobevibart (mAb) + elebsiran (siRNA)	SOLSTICE: 72 & (partial) 96-week data	✓ Jan'26
		ECLIPSE 1: topline data	4Q'26
		ECLIPSE 2: topline data	1Q'27
		ECLIPSE 3: topline data	1Q'27
PSMA-Expressing Prostate Cancer	VIR-5500 dual-masked PSMAxCD3 TCE	Phase 1 dose escalation response data	✓ Feb'26 ASCO GU
HER2-Expressing Solid Tumors	VIR-5818 dual-masked HER2xCD3 TCE	Phase 1 dose escalation response data	2H'26
EGFR-Expressing Solid Tumors	VIR-5525 dual-masked EGFRxCD3 TCE	Phase 1 initial dose escalation clinical data	TBA

CD3: cluster of differentiation 3; EGFR, epidermal growth factor receptor; HER2: human epidermal growth factor receptor 2; PSMA: prostate-specific membrane antigen; siRNA: small interfering RNA; TBA, to be announced; TCE: T-cell engager; mAb, monoclonal antibody

Our clinical programs address large and growing unmet needs

Infectious Disease

CHD

Tobevibart + elebsiran
Phase 3
Active Viremic Patients

174K

U.S.¹ + UK + EU² (all 27 member states)

Oncology – Solid Tumors

2032 prevalence estimate for U.S., EU4 and UK

PSMA

VIR-5500
Phase 1
Drug-Treated Patients³

100K
mCRPC

60K
mHSPC

HER2

VIR-5818
Phase 1
Drug-Treated Patients³

27K
HER2+ mUC

11K
HER2+ mCRC

EGFR

VIR-5525
Phase 1
Drug-Treated Patients³

431K
mNSCLC

69K
mHNSCC

271K
mCRC

¹ U.S. sources include Wong 2024, Polaris 2024, Stockdale 2020, Gish 2024

² EU sources include Polaris 2024, Delmas 2014, Wong 2024, Heidrich 2009, Reinheimer 2012, Stockdale 2020, Stroffolini 2020, Brancaccio 2019, Annual England Sentinel System 2020, Tseneva-Damyanova 2023, Papatheodoridis 2023, Parames 2016, Genne 2011, Hirzel 2015

³ Clarivate DRG, projected drug treated patients, 2032

CHD: chronic hepatitis delta; EGFR: epidermal growth factor receptor; EU4: France, Germany, Italy and Spain; HER2: human epidermal growth factor receptor 2; mCRC: metastatic colorectal cancer; mCRPC: metastatic castrate-resistant prostate cancer; mHNSCC: metastatic head and neck squamous cell carcinoma; mHSPC: metastatic hormone-sensitive prostate cancer; mNSCLC: metastatic non-small cell lung cancer; mUC: metastatic urothelial carcinoma; PSMA: prostate-specific membrane antigen

Our path to delivering transformational therapies to people living with devastating diseases: chronic hepatitis delta (CHD)



Commercializing our **chronic hepatitis delta (CHD)** combination therapy will drive near-term revenue sustainability



Accelerating our **masked T-cell engager (TCE) immunotherapy** portfolio offers key value inflection points



Aiming Vir Bio's discovery engine at **developing a robust pipeline** of cancer immunotherapies creates sustainable long-term growth




Strategic Collaborations

Selectively partner drug candidates to focus internal resources, unlock the value of our pipeline and maximize benefit to patients

CHD: devastating liver disease, significantly underserved with high mortality

CHD



¹ Negro F. (2023). Hepatitis D: A Review. *JAMA*. 330(24):2376–2387; ² Pan C, (2023). Diagnosis and Management of Hepatitis Delta Virus Infection. *Dig Dis Sci*. Aug;68(8):3237-3248; ³ Sagnelli C, et al. (2021) HBV/HDV Co-Infection: Epidemiological and Clinical Changes, Recent Knowledge and Future Challenges. *Life*,11(2):169. <https://doi.org/10.3390/life11020169>; ⁴ U.S. sources include Wong 2024, Polaris 2024, Stockdale 2020, Gish 2024; ⁵ EU sources include Polaris 2024, Delmas 2014, Wong 2024, Heidrich 2009, Reinheimer 2012, Stockdale 2020, Stroffolini 2020, Brancaccio 2019, Annual England Sentinel System 2020, Tseneva-Damyanova 2023, Papatheodoridis 2023, Parames 2016, Genne 2011, Hirzel 2015; ⁶ Stockdale A, et al. (2020). The global prevalence of hepatitis D virus infection: Systematic review and meta-analysis. *J Hepatol*, 73, 523-32
CHD: chronic hepatitis delta; HBV: hepatitis B virus; HCC: hepatocellular carcinoma
2026 Vir Biotechnology, Inc.™

Accelerating access to our CHD regimen to patients in Europe and ANZ through collaboration with Norgine

CHD




- Norgine is a leading European-focused specialty pharma with market-leading products in hepatology/GI, rare disease and pediatric oncology
- Exclusive commercial license in Europe, Australia, New Zealand
 - €55M initial reimbursement paid at closing
 - Up to €495M in clinical, regulatory and sales milestones
 - Tiered mid-teen to high-twenties percent royalties on net sales
 - ~25% sharing of future ECLIPSE external clinical costs
- Vir Biotechnology retains all commercialization rights in the U.S. and all other markets outside of the Greater China Territory¹

¹ Bii Biosciences retains rights to the combination of tobevibart and elebsiran in the Greater China Territory (People's Republic of China, Hong Kong, Taiwan and Macau)
Norgine holds exclusive license for the commercial rights to the combination of tobevibart and elebsiran in Europe, Australia and New Zealand
CHD: chronic hepatitis delta

Our CHD combination regimen is potentially best-in-class

Two complementary MOAs & highly differentiated target product profile



cccDNA: covalently closed circular DNA; CHD: chronic hepatitis delta; HBsAg: hepatitis B virus surface antigen; HBV: hepatitis B virus; HDV: hepatitis D virus; Int: integrated; MOA: mechanism of action; NRTI: nucleoside/nucleotide reverse transcriptase inhibitor; RNP: ribonucleoprotein; SVP: subviral particle


Ph2 SOLSTICE trial evaluating potential best-in-class therapy for CHD

CHD

Study design: tobevibart + elebsiran Q4W and tobevibart Q2W

Inclusion criteria:

- HDV RNA ≥ 500 IU/mL
- ALT $> \text{ULN}$; ALT $< 5 \times \text{ULN}$
- Non-cirrhotic^a or cirrhotic (CTP-A)^b
- N = 65, randomized 1:1



Primary Endpoints:

- Proportion of participants with HDV RNA $< \text{LOD}$ or $\geq 2 \log_{10}$ IU/mL reduction (virologic response) and ALT $< \text{ULN}$ (ALT response) at Week 24
- TEAEs and serious TEAEs

^a Non-cirrhotic: liver biopsy with METAVIR F0 to F3 or liver stiffness < 12 kPa within 12 months of screening and platelet count $> 150 \times 10^3/\mu\text{L}$

^b Compensated cirrhotic participants: liver biopsy with METAVIR F4 or liver stiffness ≥ 12 kPa within 12 months of screening, a platelet count $> 90 \times 10^3/\mu\text{L}$, and a CTP score of 5 or 6, inclusive at screening and at the start of the study


ALT: alanine aminotransferase; CHD: chronic hepatitis delta; CTP: Child-Turcotte-Pugh; EOT: end of treatment; HDV: hepatitis D virus; LOD: limit of detection; Q2W: once every 2 weeks; Q4W: once every 4 weeks; SC: subcutaneous; TEAE: treatment-emergent adverse event; ULN: upper limit of normal

SOLSTICE ClinicalTrials.gov Identifier: NCT05461170

Ph2 SOLSTICE trial evaluating potential best-in-class therapy for CHD

CHD

Monthly combo therapy achieved undetectable HDV RNA in 88% of patients that reached Week 96 vs. 46% with monotherapy



“
Undetectable HDV RNA is a known driver of improved CHD patient outcomes. Data from the ongoing SOLSTICE Phase 2 trial are encouraging, as they continue to show the potential of the tobevibart and elebsiran combination to achieve robust HDV suppression by tackling the viral cycle through multiple mechanisms.
”

Tarik Asselah, M.D., Ph.D.
Professor of Hepatology at the Hôpital Beaujon, APHP

CHD: chronic hepatitis delta; HDV, hepatitis D virus; mAb: monoclonal antibodies; Q2W: once every 2 weeks; Q4W: once every 4 weeks; siRNA: small interfering RNA; TND: target not detected
HDV RNA TND = undetectable HDV RNA

Data are reported for participants who completed the visit with non-missing HDV RNA and ALT or discontinued treatment before the visit

By week 96, N=3 participants discontinued tobevibart+elebsiran Q4W and N=7 participants discontinued in tobevibart Q2W and are counted as failures in analysis; n=7 and n=8


Respectively have not yet reached the week 96 visits but remain on treatment; 1 participant receiving tobevibart Q2W is censored after week 52 due to a site protocol deviation

Data as of 11/19/25

Ph2 SOLSTICE trial evaluating potential best-in-class therapy for CHD

CHD

~90% of participants receiving tobevibart + elebsiran achieved very low HBsAg values by Week 24 and maintain suppression



Hepatitis delta virus (HDV) requires serum HBV surface antigen (HBsAg) to replicate and complete its lifecycle; clearing HBsAg limits HDV replication


CHD: chronic hepatitis delta; HDV: hepatitis D virus; mAb: monoclonal antibodies; Q2W: once every 2 weeks; Q4W: once every 4 weeks; siRNA: small interfering RNA

Data are reported for participants who completed the visit with non-missing HBsAg measurement or discontinued treatment before the visit

By week 96, N=3 participants discontinued tobevibart+elebsiran Q4W and N=7 participants discontinued in tobevibart Q2W and are counted as failures in analysis; n=7 and n=8 respectively have not yet reached the week 96 visits but remain on treatment; 1 participant receiving tobevibart Q2W is censored after week 52 due to a site protocol deviation
Data as of 11/19/25

Ph2 SOLSTICE trial evaluating potential best-in-class therapy for CHD

ALT normalization was similar between tobevibart + elebsiran and tobevibart monotherapy



ALT: alanine aminotransferase; Q2W: once every 2 weeks; Q4W: once every 4 weeks; ULN: upper limit of normal

ALT ULN (male) = 40 IU/mL; ALT ULN (female) = 33 IU/mL.

Data are reported for participants who completed the visit with non-missing HDV RNA and ALT or discontinued treatment before the visit.

By week 96, N=3 participants discontinued tobevibart+elebsiran Q4W and N=7 participants discontinued in tobevibart Q2W and are counted as failures in analysis; n=7 and n=8

respectively have not yet reached the week 96 visits but remain on treatment; 1 participant receiving tobevibart Q2W is censored after week 52 due to a site protocol deviation

Data as of 11/19/25

Ph2 SOLSTICE trial evaluating potential best-in-class therapy for CHD

Majority of adverse events were grade 1 or 2 transient through Week 72

Safety or tolerability measure, n (%) ^a	Tobevibart + elebisan Q4W <i>de novo</i> N = 32	Tobevibart Q2W N = 33
Any TEAE	28 (88)	32 (97)
Grade 1-2	27 (84)	30 (91)
Grade 3	1 (3) ^b	1 (3) ^c
Grade 4	0	1 (3) ^d
Treatment-related TEAE	23 (72)	26 (79)
TEAE leading to study drug interruption	0	1 (3) ^e
TEAE leading to study drug discontinuation	0	3 (9) ^f
Serious TEAE	1 (3) ^g	1 (3) ^c
Treatment-related serious TEAE	0	0

Q2W: once every 2 weeks; Q4W: once every 4 weeks; SAE: serious adverse event; TEAE: treatment-emergent adverse event.

^aA participant with multiple events within a category is counted only once in that category.

^bGrade 3 worsening of diabetes mellitus type 2 deemed unrelated to study drugs by investigator.

^cGrade 3 hepatocellular carcinoma (SAE) deemed unrelated to study drugs by investigator.

^dGrade 4 neutropenia on Week 12 and Week 16; recovered to grade 2 or 3 after Week 16 without treatment.

^eReason for study drug interruption: neutropenia (Preferred term).

^fReason for discontinuation: 2 cases of influenza-like illness (Preferred term) and 1 case of hepatocellular carcinoma.

^gOne participants with 2 concurrent SAEs of worsening nasal septum deviation and worsening nasal concha hyperplasia (both Grade 2) who underwent planned admission to the hospital for surgery; both SAEs were deemed unrelated to study drugs by investigator.

Most TEAEs were grade 1 or 2 across treatment groups, and the most common TEAE (influenza-like illness) was generally mild to moderate and transient

Ph2 SOLSTICE results to-date show monthly tobevibart + elebsiran combo is well tolerated with robust and durable efficacy

Summary of available data through Week 96

CHD

Monthly combination therapy achieves and maintains HDV RNA TND in 88% of participants who reached Week 96

High reductions in serum HBsAg; ~90% of participants on combination therapy achieved HBsAg reductions to <10 IU/mL by Week 24 and maintained suppression

ALT normalization at Week 48 was similar between combination and monotherapy cohorts and remained stable

No grade 3 or higher treatment-related adverse events (TRAEs) with the combination therapy, and TRAEs were generally mild to moderate and transient

ALT: alanine aminotransferase; CHD: chronic hepatitis delta; HBsAg: hepatitis B surface antigen; HDV: hepatitis D virus; TND: target not detected
HDV RNA TND = undetectable HDV RNA

Registrational ECLIPSE program progressing ahead of schedule

Initial topline data anticipated in Q4 2026

CHD

✓ FDA breakthrough designation

✓ FDA Fast Track

✓ EMA PRIME designation

✓ EMA Orphan Drug designation

ECLIPSE 1

Phase 3

- HDV RNA TND + ALT normalization at **Week 48**
- Tobeivart + elebsiran vs. deferred treatment (n=120, 2:1)



Fully enrolled

ECLIPSE 2

Phase 3

- HDV RNA TND at **Week 24**
- Tobeivart + elebsiran vs. bulevirtide switch* (n=150, 2:1)

Enrollment
On Track

ECLIPSE 3

Phase 2b

- HDV RNA TND at **Week 48**
- Tobeivart + elebsiran vs. bulevirtide naïve (n=100, 2:1)



Fully enrolled

*Defined as failure to achieve HDV RNA < 500 IU/mL with bulevirtide
ALT: alanine aminotransferase; HDV: hepatitis D virus; TND: target not detected
HDV RNA TND = undetectable HDV RNA
ECLIPSE ClinicalTrials.gov Identifiers: ECLIPSE 1 NCT06903338, ECLIPSE 2 NCT07128550, ECLIPSE 3 NCT07142811

Our path to delivering transformational therapies to people living with devastating diseases: cancer immunotherapy



Commercializing our **chronic hepatitis delta (CHD)** combination therapy will drive near-term revenue sustainability



Accelerating our **masked T-cell engager (TCE) immunotherapy** portfolio offers key value inflection points



Aiming Vir Bio's discovery engine at **developing a robust pipeline** of cancer immunotherapies creates sustainable long-term growth



Strategic Collaborations

Selectively partner drug candidates to focus internal resources, unlock the value of our pipeline and maximize benefit to patients

T-cell engagers (TCEs) are a powerful modality in cancer therapy

Our masked TCEs act like Trojan Horses, powered by the PRO-XTEN® platform

TCEs hold tremendous potential, limited by toxicity

- 10 TCE breakthrough immunotherapies already on the market¹
- Application in solid tumors limited due to toxicity and off-tumor activation
- Masking ensures TCEs are **only activated in the tumor microenvironment**

The PRO-XTEN® masking platform

Clinically validated, used on a blockbuster drug for hemophilia A²

Universal, plug-and-play platform enables acceleration of next generation of drug candidates

Our masked TCEs act like Trojan Horses, designed to maximize therapeutic index

Masks cleaved off by the proteases in the tumor microenvironment

Designed to reduce toxicity, enabling higher dosing and **wider therapeutic window**

Longer drug half-life supports **optimization of dosing schedules**

¹ Glaser, A., Kochanowski, K., Oh, D., Porritt, R. A., & Kim, H. (2024). T cell engagers emerge as a compelling therapeutic strategy for solid tumors. Journal of Experimental Medicine

² ALTUVIIIIO® [Antihemophilic Factor (Recombinant), Fc-VWF-XTEN Fusion Protein-ehtl] is marketed for hemophilia A and is a registered trademark of Sanofi



Q3W: once every three weeks

Our unique pipeline of TCEs is enabled by the PRO-XTEN® masking platform

Allows our TCEs to overcome challenges of unmasked and single-masked TCEs


TCE

Shields the core of TCEs, expanding potential in cancer therapy



Tumor-binding domain

Variable region binds tumor-associated antigen



T-cell-binding domain

Variable region binds CD3 to recruit T-cells



PRO-XTEN® mask

XTEN masks off-tumor activity of the TCE and prolongs half-life



Cleavable linkers

Proteases in the TME selectively cleave linkers to release masks



Phase 1 Clinical Data: **VIR-5500 (PSMA)**

Presented February 2025

Landmark strategic collaboration with Astellas maximizes potential of VIR-5500 in prostate cancer

VIR-5500
(PSMA)



Collaboration pairs Astellas' world class capabilities in prostate cancer with Vir Bio's potential best-in-class T-cell engager VIR-5500, powered by PRO-XTEN[®] masking technology

\$1.7B

in upfront payments and milestones¹

50/50

profit/loss share in the U.S.
Co-promote option for
Vir Bio

40/60 Vir Bio / Astellas global
development cost share²
Tiered, **double-digit**
royalties on ex-U.S. net sales

Positions Vir Bio to accelerate clinical development of VIR-5500 into pivotal trials in 2027

¹ Amounts shown exclude payments to third parties. Sanofi is entitled to 20% of certain collaboration proceeds, including: upfront, equity premium, and the portion of milestones, profit share & royalties that exceed the amounts already owed to Sanofi under the terms of the existing Sanofi agreement, effective September 9, 2024.

² R&D cost share: Global studies Vir Bio 40% & Astellas 60%; U.S.-specific studies Vir Bio 50% & Astellas 50%; ex-U.S.-specific studies Astellas 100%

VIR-5500 monotherapy study of PSMA-targeted dual-masked TCE in prostate cancer

All dose escalation cohorts have cleared DLT (N=58)

VIR-5500
(PSMA)

QW dose escalation (N=26)

1000 → 2000 → 3000 µg/kg (N=4)

500 → 1000 → 2000 µg/kg (N=3)

300 → 600 → 1000 µg/kg (N=4)

200 → 300 → 400 µg/kg (N=5)

120 → 180 → 180 µg/kg (N=4)



60 µg/kg (N=3)



30 µg/kg (N=3)

Q3W dose escalation (N=32)

1000 → 2000 → 4000 µg/kg
w/ Prophylactic Steroids (N=3)

800 → 2000 → 3500 µg/kg (n=9)

800 → 1500 → 3000 µg/kg (N=5)

1000 → 2000 → 3000 µg/kg (N=5)



500 → 1000 → 2000 µg/kg (N=4)

300 → 600 → 1000 µg/kg (N=6)

Eligibility criteria:

- Documented progressive metastatic CRPC
- ≥ 1 prior taxane regimen
- ≥ 1 prior ARPI
- 0 to 1 ECOG status
- Life expectancy >6 months

No requirement of prophylactic steroids or IL-6 therapy except for exploratory analysis in high dose cohort (n=3)

ARPI: androgen receptor pathway inhibitor; CRPC: castration-resistant prostate cancer; DLT: dose limiting toxicities; ECOG: Eastern Cooperative Oncology Group; PSMA: prostate-specific membrane antigen; QW: once weekly; Q3W: once every 3 weeks; TCE: T-Cell Engager
VIR-5500-V101 ClinicalTrials.gov Identifier: NCT05356741 Data as of: January 9, 2026

Late-line mCRPC population including patients with liver metastases

VIR-5500
(PSMA)

VIR-5500 baseline characteristics NCT05997615	VIR-5500 N=58
Median age, years (range)	68 (49-81)
Prior lines of therapy (Any Setting)	
Number, Median (Min, Max)	4 (2, 7)
Prior Taxane, n (%)	55 (94.8)
Prior ARPI, n (%)	58 (100)
Prior PSMA-radioligand therapy ^a , n (%)	7 (12.1)
Baseline Central PSA, ng/mL	79
Median (min, max)	(4, 3708)
Disease Characteristics	
RECIST-evaluable ^b , n (%)	30 (51.7)
Bone metastases, n (%)	52 (92.9)
Lymph node metastases, n (%)	18 (32.1)
Visceral metastases ^c , n (%)	25 (44.6)
Liver metastases, n (%)	10 (17.9)

Study & enrollment details:

Heavily pre-treated participants:

- Median 4 prior lines of therapy
- 95% prior taxanes in any setting*
- 12% prior PSMA-radioligand therapy (mostly in low dose cohorts)
- 2 with prior STEAP1 TCE

Significant disease burden in all cohorts:

- 93% of subjects with bone metastases
- 45% visceral metastases
 - 18% liver metastases (poor prognosis)

*Participants who were deemed clinically unsuitable to be treated with a taxane regimen or have refused treatment with a taxane regimen are considered eligible.

^a 4 of 7 Prior PSMA-radioligand therapy treated patients received VIR-5500 doses ≤ 120 μ g/kg.

^b RECIST-evaluable population is defined as having measurable disease documented at baseline according to RECIST v1.1 criteria and at least one follow-up tumor assessment after a full cycle of treatment

^c Visceral metastases include site of lesions in lung, adrenal and liver.

Max: maximum; mCRPC; metastatic castration-resistant prostate cancer; Min: minimum; N: number of participants; PSA: prostate-specific antigen; PSMA: prostate-specific membrane antigen; RECIST: Response Evaluation Criteria in Solid Tumors; STEAP1: Six-transmembrane epithelial antigen of the prostate 1; TCE: T Cell Engager

VIR-5500-V101 ClinicalTrials.gov Identifier: NCT05997615. Data as of: January 9, 2026

VIR-5500 Phase 1 study shows favorable safety profile with meaningful anti-tumor activity

VIR-5500
(PSMA)

Well tolerated with no dose limiting toxicities

- 12% Grade ≥ 3 TRAEs
- Limited CRS 50% (29/58) primarily seen in first cycle and mostly Grade 1 (Fever Only)
- No DLTs

Dose-dependent and meaningful anti-tumor activity

- Clear dose-response relationship with deeper PSA declines observed with higher doses
- 45% (5/11) ORR of RECIST-evaluable* patients (Q3W $\geq 3,000$ $\mu\text{g/kg}$)
 - 4 patients confirmed up to Week 27
 - 1 patient pending confirmation

Early signs of PSA and radiographic durability

- Emerging evidence of durable PSA₅₀, PSA₉₀ and RECIST responses

$\geq 3,000$ $\mu\text{g/kg}$ Q3W

CRS

All Grades	59% (13/22)
Grade 1	50% (11/22)
Grade 2	9% (2/22)
Grade 3	0

PSA Response

PSA ₅₀	82% (14/17)
PSA ₉₀	53% (9/17)
PSA ₉₉	29% (5/17)

Safety evaluable population is defined as all enrolled participants who have received at least one dose of VIR-5500.

PSA evaluable population is defined as participants who received ≥ 1 cycle of VIR-5500 and must have at least one pre-treatment and at least one post-treatment PSA measurement.


PSA₅₀, PSA decline of 50%-100% from baseline;
PSA₉₀, PSA decline of 90%-100% from baseline;
PSA₉₉, PSA decline of 99%-100% from baseline

*RECIST-evaluable population is defined as having measurable disease documented at baseline according to RECIST v1.1 criteria and at least one follow-up tumor assessment after a full cycle of treatment
CRS: cytokine release syndrome; DLT: dose limiting toxicities; G3+: Grade ≥ 3 ; ORR: objective response rate; PSA: prostate-specific antigen; Q3W: once every 3 weeks; RECIST: Response Evaluation Criteria in Solid Tumors; TRAE: treatment related adverse event
VIR-5500-V101 ClinicalTrials.gov Identifier: NCT05997615. Data as of: January 9, 2026

Case Study 1: complete resolution of multiple (>14) liver lesions at Week 9; PSA₉₉

VIR-5500
(PSMA)

Q3W Cohort 800/1500/3000 µg/kg




Case study detail

- 63-year-old male
- High-disease burden: liver and bone lesions
- 5 Prior Lines of Treatment: Docetaxel, Olaparib, Cabazitaxel, Abiraterone, MOMA-313-001
- **uPR, 63% decrease in tumor diameter**
- Metabolic response of PSMA-avid bone and hepatic lesions
- Patient bone pain resolved
- Continues on treatment (Cycle 6)

Case Study 2: significant RECIST response in large liver lesions

VIR-5500
(PSMA)

Q3W cohort 800/2000/3500 µg/kg




Case study detail

- 75-year-old male
- High-disease burden; liver and lymph-nodes (no bone lesions)
- 6 Prior Lines of Treatment: Enzalutamide, Docetaxel, Cabazitaxel, and NX-1607
- **Confirmed PR, 46% decrease in tumor diameter**
- Disappearance of majority of PSMA-avid hepatic lesions
- Continues on study (Cycle 10)

Case Study 3: durable RECIST, PSMA-PET and deep PSA response up to 8 months

VIR-5500
(PSMA)



Q3W cohort 800/2000/3500 µg/kg



Case study detail

- 70-year-old male with peritoneal and abdominal wall lesions
- 3 Prior lines of Treatment: Enzalutamide, Docetaxel, Cabazitaxel
- **Confirmed PR**, complete resolution of small lesions
- Ongoing sustained PSA₉₀ response at 8 months
- Complete PSMA-PET response
- Excellent Quality of life
- Continues on study (Cycle 10)

Case Study 4: complete response in diffuse lesions with prior radioligand therapy




Case Study Detail

- 63-year-old male with lymph-node and bone lesions
- 5 prior lines of Treatment including Enzalutamide, Docetaxel, TAS3681 AR antagonist, and RLT (²²⁵Ac-pelgifatama)
- **CR for target lesions** and non-CR/non-PD for non-target bone lesions; PSA₉₉ at Cycle 2 Day 1 (Week 4)
- Excellent quality of life, with significant reduction in pain
- Patient withdrew from the study on Cycle 5, while in remission since deriving significant benefit

Case Study 5: complete response with ~12 months of durability

VIR-5500
(PSMA)




Case study detail

- 77-year-old male
- Significant disease burden including >20 bone lesions and positive lymph nodes
- 4 prior lines of Treatment:
- Darolutamide, Abiraterone, Olaparib, and Docetaxel
- **Complete Response** per PCWG3 at Cycle 9/Week 27
- Significant PSA responder with PSA₉₀ response starting at Cycle 3 Day 1 and PSA currently undetectable
- Continues on treatment (Cycle 17)

Q3W cohort 300/600/1000 µg/kg




Baseline

PSA 42 ng/mL



Week 9

PSA 0.6 ng/mL



Baseline Week 9 Week 18 Week 27

Well-tolerated with favorable safety profile

12% grade ≥ 3 TRAEs, limited CRS, mostly grade 1 (Fever Only), and no DLTs

Limited High-Grade Events and Tx Discontinuations

TEAEs in any Participant n (%) (N=58)

Any TEAE	58 (100)
Related TEAE	50 (86.2)
Serious Related TEAE	17 (29.3)
Related Grade ≥ 3 TEAE [#]	7 (12.1)
TEAE Leading to Treatment Discontinuation [^]	2 (3.4)

[#] Related Grade ≥ 3 TEAE: AST increased, neutrophil count decreased, WBC count decreased, cytokine release syndrome (in participant at 200-300-400 ug/kg undergoing intraparticipant dose escalation), tumor flare/arthritis, neutropenia, blurred vision, lymphocyte count decreased, bone pain


[^] One unrelated spinal cord compression requiring radiation therapy and one Grade 4 treatment-related blurred vision event that improved, with unclear pathophysiology & non-specific MRI findings.

Treatment Related AEs (Most Frequent) in Doses $\geq 3,000$ $\mu\text{g/kg}$ Q3W (N=22)

Preferred Term	Grade 1	Grade 2	Grade ≥ 3
Participant with at least 1 TRAE	9 (41)	9 (41)	3(14) ¹
Cytokine Release Syndrome	11 (50)	2 (9)	0
Back Pain	3 (14)	2 (9)	0
Fatigue	3 (14)	2 (9)	0
Infusion Related Reaction	4 (18)	1 (5)	0
Anemia	1 (5)	3 (14)	0
Asthenia	4 (18)	0	0
Nausea	4 (18)	0	0
Blurred Vision	1 (5)	0	2 (9)

¹ G3 Neutropenia (2), G3 Tumor Flare, G3 Bone Pain (2), G3 Lymphocyte Decrease; G3 Vision Blurred, G4 Vision Blurred

Low Grade CRS in Doses $\geq 3,000$ $\mu\text{g/kg}$ Q3W (N=22)



✓ No DLTs*

✓ No ICANS

✓ Limited transaminase elevation

✓ Very Low incidence of hypoacusis, xerostomia, dry eye, stomatitis or dysgeusia; Grade 1 only¹

No requirement of prophylactic steroids or IL-6 therapy except for exploratory analysis in high dose cohort (n=3)

* Two blurred vision events at the 4,000 ug/kg dose were reviewed and incorporated into dose-escalation considerations and ongoing study conduct

[^] 4-8 mg of Dexamethasone premedication in cycle 1 evaluated in 1000-2000-4000 ug/kg Q3W cohort.


¹ Grade 1 events: Dry Mouth/Xerostomia (4), Dry Eye (2), Hypoacusis (5), Stomatitis (1), Dysgeusia (1), Renal toxicities (2); as well as 3 unrelated SAEs of Acute Kidney Injury and Hematuria

AE: adverse event; C1: cycle one; CRS: cytokine release syndrome; DLT: dose limiting toxicities; ICANS: immune effector cell-associated neurotoxicity syndrome; Q3W: once every 3 weeks; TEAE: treatment emergent adverse event

VIR-5500-V101 ClinicalTrials.gov Identifier: NCT05997615. Data as of: January 9, 2026

Dose response relationship: deeper PSA declines and confirmed PSA responses at higher doses


VIR-5500
(PSMA)



Deep PSA declines observed as early as Cycle 1 Day 8, evidence of concordant RECIST responses in evaluable patients

VIR-5500
(PSMA)

Doses ≥ 3000 $\mu\text{g/kg}$ Q3W



≥ 3000 $\mu\text{g/kg}$ Q3W

Any PSA Decline	15/17 (88%)
PSA ₅₀	14/17 (82%)
PSA ₉₀	9/17 (53%)
PSA ₉₉	5/17 (29%)


Significant Anti-tumor Responses:

- Rapid and deep responses as soon as Cycle 1 Day 8
- Participants with the deepest PSA responses (PSA₉₀ & PSA₉₉) often had confirmed RECIST responses

PSA₅₀, PSA decline of 50%-100% from baseline; PSA₉₀, PSA decline of 90%-100% from baseline; PSA₉₉, PSA decline of 99%-100% from baseline. Evaluable participants who received ≥ 1 cycle of VIR-5500. PCWG3 measured PSA at $\geq 2\text{D1}$. Y-axis truncated at 100. VIR-5500-V101 ClinicalTrials.gov Identifier: NCT05997615. Data as of: January 9, 2026

Reduction in PSMA total tumor volume by central PSMA PET analysis, RECIP responses associated with deep PSA declines and RECIST responses

Doses ≥ 3000 ug/kg Q3W



Imaging performed at baseline and at 9 weeks, or EOT, whichever comes first.
PSMA-Total Tumor Volume is the sum of all PSMA-avid lesions (bone and soft tissue) in cubic centimeters.
<https://pubmed.ncbi.nlm.nih.gov/40473460/>
Y-axis truncated at 100%


C1: cycle one; cCR: confirmed complete response; cPR: confirmed partial response; EOT: end of treatment; NM: non-measurable disease; PD: progressive disease; PET: positron emission tomography; PSA: prostate-specific antigen; PSMA: prostate-specific membrane antigen; Q3W: once every 3 weeks; RECIP: Response Evaluation Criteria in PSMA; RECIST: Response Evaluation Criteria in Solid Tumors; SD: stable disease; uPR: unconfirmed partial response

¹ Prior Radioligand therapy

Robust RECIST responses with VIR-5500 monotherapy in RECIST-evaluable participants

VIR-5500
(PSMA)

Doses ≥ 3000 $\mu\text{g/kg}$ Q3W



Measurable Lesion Site	Liver	Liver, Adrenal	Adrenal gland	Liver, Abdom wall	Lymph nodes	Lymph node	Liver, LN	Lymph Nodes	Peritoneal implants	Lymph node	Liver
Best Overall Response	PD	PD	PD (new lesions)	SD	SD	PD (new lesions)	cPR	cPR	cPR	cCR	uPR (Wk 18 pending)
PSMA PET Response (SUVmax)	Not available	-79.6%	8.3	-78%	-64.1%	-22.6%	-89.7%	-95%	-81%	-88.2%	-97.3%

¹ Prior Radioligand therapy

* 4 patients with confirmed responses and 1 patient pending confirmation

Imaging performed every 9 weeks

Baseline Assessment: patient must have measurable disease per RECIST criteria at baseline


Post-Baseline Assessment: at least one follow-up tumor assessment after starting treatment is required to determine response (CR, PR, SD, PD)

C1: cycle one; cCR: confirmed complete response; cPR: confirmed partial response; DCR: Disease Control Rate (includes SD+PR+CR); NM: non-measurable disease; ORR: objective response rate; PD: progressive disease; PET: positron emission tomography; PR: partial response; PSA: prostate-specific antigen; PSMA: prostate-specific membrane antigen; Q3W: once every 3 weeks; RECIST: Response Evaluation Criteria in Solid Tumors; SD: stable disease; SLD: sum of longest diameters; QW: once weekly; PD: progressive disease; uPR: unconfirmed partial response

RECIST responses concordant with deep PSA responses

VIR-5500
(PSMA)

Doses ≥ 3000 $\mu\text{g/kg}$ Q3W



Radiographic responses at efficacious Q3W dose levels

- 45% (5/11) ORR in RECIST-evaluable patients
- All evaluable responders achieved confirmed responses; 1 patient pending confirmation
- Tumor shrinkage across multiple lesions and patients, providing evidence of effective T-cell engagement

Imaging performed every 9 weeks

* Week18 visit pending


C1: cycle one; cCR: confirmed complete response; cPR: confirmed partial response; ORR: objective response rate; PSA: prostate-specific antigen; PSMA: prostate-specific membrane antigen; Q3W: once every 3 weeks; RECIST: Response Evaluation Criteria in Solid Tumors; SLD: sum of longest diameters; uPR: unconfirmed partial response

VIR-5500-V101 ClinicalTrials.gov Identifier: NCT05997615. Data as of: January 9, 2026

Emerging evidence of PSA and radiographic durability

VIR-5500
(PSMA)

Doses ≥ 3000 $\mu\text{g/kg}$ Q3W, CRS Restricted to Early Cycles



*101-1007: PSMA-positive brain lesions, early progressor, received 3 cycles of VIR-5500

^101-1006: PSMA-negative liver lesions, early progressor, received 2 cycles of VIR-5500

cCR: confirmed complete response; cPR: confirmed partial response; CRS: cytokine release syndrome; G1: Grade 1; G2: Grade 2;

PSA: prostate-specific antigen; Q3W: once every 3 weeks; uPR: unconfirmed partial response

VIR-5500-V101 ClinicalTrials.gov Identifier: NCT05997615. Data as of: January 9, 2026

Next steps for VIR-5500: rapid advancement into earlier treatment setting in prostate cancer

VIR-5500
(PSMA)

Conclusion of monotherapy QW and Q3W dose escalation in late-line mCRPC

Initial monotherapy recommended expansion dose

Progressing toward initiating Expansion Dose Cohorts in 2Q'26

- Late-line mCRPC (Monotherapy)
- Early-line mCRPC (Combination)
- mHSPC (Combination)

Plans to initiate Phase 3 program in 2027

QW: once weekly; Q3W: once every 3 weeks; mCRPC: metastatic castration-resistant prostate cancer; mHSPC: metastatic hormone-sensitive prostate cancer

Our clinical pipeline of masked TCEs demonstrates promise of the PRO-XTEN[®] platform

TCE

VIR-5500 (PSMA)



The **only dual masked TCE** in clinical development for prostate cancer

Well-tolerated with favorable safety profile

Potent anti-tumor activity

Potential to move into earlier treatment settings and initiate pivotal trials in 2027

VIR-5818 (HER2)



The **only masked TCE** in clinical development for HER2 tumors

PD-1 combination dose escalation **ongoing**

Phase 1 dose escalation data in 2H'26

VIR-5525 (EGFR)



Dual-masked TCE in clinical development for EGFR tumors (incl. NSCLC, CRC, HNSCC and others)

First patient dosed on July 25, 2025

Phase 1 initial dose escalation data TBD

7 Preclinical Programs



Universal masking platform allows us to rapidly expand into other solid tumors with high unmet need

7 preclinical programs across solid tumors, including lung, colorectal and bladder cancers

Progressing to development candidate selection by early 2027

CRC: colorectal cancer; EGFR: epidermal growth factor receptor; HER2: human epidermal growth factor receptor 2; HNSCC: head and neck squamous cell carcinoma; NSCLC: non-small cell lung cancer; PSMA: prostate-specific membrane antigen; TCE: T-cell engager

Our strategic approach creates short and long-term value drivers



chronic hepatitis delta

ECLIPSE 1 topline data
in 4Q26

ECLIPSE 2 & 3 topline data in
1Q27



universal masked TCEs

VIR-5500 PSMA pivotal trials
to initiate in 2027

VIR-5818 HER2 data update
in 2H26



discovery engine

7 preclinical PRO-XTEN® TCE
targets identified

Progressing to development
candidate selection by early
2027



Strategic Collaborations

Selectively partner drug candidates to focus
internal resources, unlock the value of our
pipeline and maximize benefit to patients

Financial Highlights

\$782M cash and investments¹ with cash runway
into Q2 2028²

¹ Vir Bio reported cash, cash equivalents and investments of \$782 million as of December 31, 2025.

² Cash runway projection based on the current operating plan and incorporating the expected effects of the Astellas collaboration agreement.
Numbers above may not tie due to rounding.

PATIENTS ARE WAITING



Phase 1 Clinical Data: **VIR-5818 (HER2)**

Presented January 2025

The first clinical stage masked HER2 TCE in ongoing Phase 1

VIR-5818
(HER2)

Part 1: Monotherapy Dose Escalation - Completed

Recommended expansion
dose and schedule

100 → 300 → 1000 µg/kg

100 → 300 → 800 µg/kg¹

100 → 250 → 600 µg/kg

100 → 200 → 400 µg/kg

200 µg/kg

1 µg/kg

Eligibility:

HER2 IHC2-3+, ISH+, or
mutant

Exhausted all SOC

79 patients enrolled

Evaluating QW and Q3W

Demonstrates wide safety
margin

Part 2: Pembrolizumab Combination

VIR-5818
QW and Q3W



Pembrolizumab
Q3W
200 mg

Currently enrolling

Analysis ongoing

Currently Evaluating

Cleared DLT


¹ Evaluating 800 µg/kg Q3W maintenance dose (cycle ≥ 2) in parallel
HER2: human epidermal growth factor receptor 2; IHC: immunohistochemistry; ISH: in situ hybridization; QW: once weekly; Q3W: once every 3 weeks; SOC:
standard of care; TCE: T-cell engager
VIR-5818 ClinicalTrials.gov Identifier: NCT05356741; Data cutoff: November 11, 2024

Minimal unmasked TCE in circulation and potential for Q3W dosing

VIR-5818
(HER2)

Minimal unmasked TCE outside the tumor


VIR-5818 and uTCE PK, First Cycle*



Low levels of uTCE in circulation,
consistent with minimal CRS

Half-life of ~ 6 days unlocks potential Q3W dosing

VIR-5818 PK, First Dose



Linear and dose
proportional PK

Preliminary safety data indicates VIR-5818 is not dose-limited by CRS

VIR-5818
(HER2)


Highly Tolerable Safety

TRAE (max grade) in >15% of pts

VIR-5818 N = 79	Grade 1 N (%)	Grade 2 N (%)	Grade ≥ 3 N (%)
Any TRAE	15 (19.0)	35 (44.3)	13 (16.5)
Pneumonitis*	16 (20.3)	9 (11.4)	2 (2.5)*
CRS	16 (20.3)	8 (10.1)	0
Nausea	12 (15.2)	8 (10.1)	0
Asthenia	12 (15.2)	6 (7.6)	1 (1.3)
Diarrhoea	14 (17.7)	5 (6.3)	0
Pruritus	13 (16.5)	1 (1.3)	0
Vomiting	8 (10.1)	6 (7.6)	0

Low Cytokine Levels, Even at Higher Doses

Peaks of IL-6 Secretion Post VIR-5818 Dosing



IL-6 release significantly lower than for
unmasked TCEs, despite higher VIR-5818 dose

¹Wermke et al., ASCO-SITC 2018

*Two cases of G3 pneumonitis include one event reversible with treatment and one confounded by rapid progression of pulmonary disease


CRS: cytokine release syndrome; IL-6: interleukin-6; TCE: T-cell engager; TRAE, treatment-related adverse event

VIR-5818 ClinicalTrials.gov Identifier: NCT05356741; Data cutoff: November 11, 2024

Notable tumor shrinkage observed during dose escalation

HER2+ Solid Tumors

(Doses ≥ 400 $\mu\text{g/kg}$)



No of prior lines	3	7	3	4	4	4	2	3	6	5	1	3	3	1	1	4	1	4	6	9
Liver metastases																				

*Includes cPR, uPR, and mixed responses

Note: HER2+ defined as IHC3+ or ISH+

BC: breast cancer; CCA: cholangiocarcinoma; cPR: confirmed partial response; CRC: colorectal cancer; DCR: disease control rate; GEJ: gastroesophageal junction; HER2: human epidermal growth factor receptor 2; IHC: immunohistochemistry; ISH: in situ hybridization; OC: ovarian cancer; PAAD: pancreatic adenocarcinoma; QW: once weekly; Q3W: once every 3 weeks; RECIST: Response evaluation criteria in solid tumors; SCCAC: squamous cell carcinoma of the anal canal; SLD: sum of longest diameters; uPR: unconfirmed partial response; VIR-5818 ClinicalTrials.gov Identifier: NCT05356741; Data cutoff: November 11, 2024

Efficacy detail:

- ≥ 400 $\mu\text{g/kg}$ drive significant RECIST responses
 - Dose escalation continues in QW and Q3W regimens
- 50% observed tumor shrinkage (10/20 patients), with a DCR of 65%
 - 4/20 responses to date*
 - Responses in patients with up to 9 prior lines
 - 14/20 with prior HER2 treatment

A patient's journey: dramatic response in advanced HER2+ breast cancer

VIR-5818
(HER2)

Tumor pain, inflammation

Day 1 Baseline



Cycle 1 Day 8



Cycle 2 Day 1



Cycle 2 Day 8



Cycle 3 Day 8



Cycle 4 Day 1



VIR-5818 Case Study

Compelling activity in breast cancer patient by Cycle 1 with transformative clearance of tumor

9 prior lines of therapy, including Enhertu


Dose: 100/300/1000 $\mu\text{g/kg}$
Well-tolerated

52% tumor shrinkage from baseline

Deep responses at early doses in MSS colorectal cancer, a tumor type traditionally resistant to immunotherapy

HER2+ Colorectal Cancer

(All Patients Shown are MSS)



Early Phase 1 efficacy

Activity	HER2+ CRC ≥400 µg/kg
cPR	2/6 (33%)
CEA Response*	3/3 (100%)
DCR ¹	5/6 (83%)


- 33% response and 100% biomarker response in mCRC
- Up to 18.1 months duration of response (pt remains on study)
- Significant room to dose escalate; potential for Q3W dosing

Note: HER2+ defined as IHC3+ or ISH+
* CEA response defined as >50% decrease in CEA post-treatment. Denominator includes all pts with longitudinal data
¹ DCR defined as stable disease or better
CEA: carcinoembryonic antigen; cPR: confirmed partial response; CRC: colorectal cancer; DCR: disease control rate; HER2: human epidermal growth factor receptor 2; IHC: immunohistochemistry; ISH: in situ hybridization; MSS: microsatellite stability; SLD: sum of longest diameters
VIR-5818 ClinicalTrials.gov Identifier: NCT05356741; Data cutoff: November 11, 2024

Patient Case Study: 2 years on treatment, exceptional durability

VIR-5818
(HER2)


Rapid and Sustained Decrease Over Time



Rapid and sustained CEA decrease with deeper tumor shrinkage when dose escalates

- 57-Year-old male w/ colorectal cancer (MSS/TMB Low)
- Status: remains on study (current dose: 600 µg/kg QW)
- HER2 status: IHC 3+

Dose-Dependent Tumor Shrinkage




- 6 prior lines including trastuzumab / tucatinib
- Significant improvement on quality of life
- 114 doses as of data cutoff, patient remains on study

CEA: carcinoembryonic antigen; cPR: confirmed partial response; IHC: immunohistochemistry; MSS: microsatellite stability; QW: weekly; SLD: sum of longest diameters; TMB: tumor mutational burden
VIR-5818 ClinicalTrials.gov Identifier: NCT05356741; Data cutoff: November 11, 2024

Molecular evidence of anti-tumor activity across multiple cancer types

Molecular Responses: ctDNA (Step-up doses only)



Detail:

- High value of biomarkers for immunologics
- RECIST responses may be confounded by tumor inflammation
- With on-treatment ctDNA collection, VIR-5818 has **molecular response for 54% subjects¹**
- Now universally collecting ctDNA

¹ Molecular response defined as >50% decline in overall ctDNA
BC: breast cancer; CRC: colorectal cancer; ctDNA: circulating tumor DNA; GEJ: gastroesophageal junction; NSCLC: non-small-cell lung cancer;
RECIST: Response evaluation criteria in solid tumors
VIR-5818 ClinicalTrials.gov Identifier: NCT05356741; ctDNA data cutoff: November 19th, 2024; EDC data cutoff: November 11, 2024

A potential first-in-class HER2 TCE designed to clinically validate the PRO-XTEN[®] platform

VIR-5818
(HER2)

Clear activity based on early Phase 1 data with potential for long-term durable responses

Emerging activity: wide TI in heavily pretreated population

- Unprecedented tolerability: no Gr3+ CRS, 16% all Gr3+ TRAEs
- 33% response in heavily pre-treated CRC patients (≥ 400 $\mu\text{g/kg}$)
- ctDNA Molecular response in 54% of subjects

Proof of concept for PRO-XTEN[®] platform

- Clear evidence of unmasking with antitumor activity

Universal masks: mechanism designed to apply across platform

- Potential rapid dose escalation for VIR-5500 (PSMA) and other targets

CRC: colorectal cancer; CRS: cytokine release syndrome; ctDNA: circulating tumor DNA; Gr3: Grade 3; HER2: human epidermal growth factor receptor 2; PSMA: prostate-specific membrane antigen; TCE: T-cell engager; TI: therapeutic index; TRAE: treatment-related adverse event
VIR-5818 ClinicalTrials.gov Identifier: NCT05356741; Data cutoff: November 11, 2024




Phase 1 Clinical Program: **VIR-5525 (EGFR)**


VIR-5525 Phase 1 study design: dose escalation and expansion

VIR-5525
(EGFR)


Part 1 & 2: Monotherapy Dose Escalation & Expansion



Part 3 & 4: Pembrolizumab Combination Dose Escalation & Expansion




Note: Step up dosing and additional schedules may be evaluated based on emerging clinical and PK data




Preclinical data demonstrate potent activity and substantial safety margin with PRO-XTEN[®] masking

VIR-5525
(EGFR)

Outside Tumor: masking is maintained, leading to
~10,000-fold shift in cytotoxicity



In Tumor: similar anti-tumor activity in
PRO-XTEN[®] masked vs. unmasked EGFR TCE



PRO-XTEN[®] masked EGFR TCE enabled ~250-fold higher tolerated exposure in NHPs vs. unmasked TCE²

¹ Cattaruzza, F., Nazeer, A., To, M. et al. Precision-activated T-cell engagers targeting HER2 or EGFR and CD3 mitigate on-target, off-tumor toxicity for immunotherapy in solid tumors. Nat Cancer 4, 485–501 (2023). <https://doi.org/10.1038/s43018-023-00536-9>. ~250-fold safety margin data from IND filing analysis (>200-fold reported in Nature paper).

² Adapted from Cattaruzza, F., Nazeer, A., Lange, Z., Hammond, M., Koski, C., Henkensiefken, A., & Schellenberger, V. (2020). HER2-XPAT and EGFR-XPAT: Pro-drug T-cell engagers (TCEs) engineered to address on-target, off-tumor toxicity with potent efficacy in vitro and in vivo and large safety margins in NHP. Cancer Research, 80(16_Supplement), 3376-3376.

EGFR: epidermal growth factor receptor; EGFR uTCE: EGFR-targeted unmasked T-cell engager; mpk: milligrams per kilogram; NHP: non-human primate; TCE: T-cell engager

© 2026 Vir Biotechnology, Inc.™